# Human bone points from Mesolithic Doggerland

## Joannes Dekker<sup>1,2,3</sup>, Virginie Sinet-Mathiot<sup>4</sup>, Merel Spithoven<sup>5</sup>, Bjørn Smit<sup>6</sup>, Arndt Wilcke<sup>7</sup>, Frido Welker<sup>2,4</sup>, Alexander Verpoorte<sup>1</sup>, Marie Soressi<sup>1</sup> <sup>1-</sup> Faculty of Archaeology<sup>,</sup> Universiteit Leiden<sup>, 2-</sup> Section for Evolutionary Genomics<sup>,</sup> Globe Institute<sup>,</sup> University of Copenhagen<sup>, 3-</sup> Department of Archaeology<sup>,</sup> University of York<sup>,</sup>

<sup>4</sup> Department of Human Evolution, Max Planck Institute for Evolutionary Anthropology, <sup>5</sup> Institute of Archaeology, University of Gronigen, <sup>6</sup> Cultural Heritage Agency, <sup>7</sup> Fraunhofer Institute for Cell Therapy and Immunology, IZI Fraunhofer,

#### Introduction

Mesolithic barbed osseous points are regularly deposited on the Dutch coast. Microwear research indicates that these points were intensively used. Due to heavy modifications the points cannot be taxonomically identified using morphology.

We applied ZooMS to investigate potential patterns of species selection in these heavily curated tools.

#### **Materials**

For our study we analysed 10 barbed points from five beaches (Fig. 1). Nine of these points are bone, the other is antler.

## Methods

All barbed points were analysed according to the Cold Acid and AmBic ZooMS protocol (Fig. 2). A database of peptide marker series for all European, Pleistocene medium to large size mammals was used for taxonomic identification (Welker et al. 2016).

Additionally the barbed points were <sup>14</sup>C dated and their  $\delta^{13}$ C and  $\delta^{15}$ N isotope ratios were measured using protocols described in Van der Plicht *et al.* 2016.

#### Acknowledgements

Prof. dr. H. van der Plicht, dr. S. Palstra for advice and assistance in sampling and for the analysis at the Centre for Isotope Research, University of Groningen, the Netherlands. Prof. dr. J.J. Hublin for support and use of facilities of the Max Planck Institute of Evolutionary Anthropology, Leipzig. E. Mulder, Prof. dr. A. van Gijn for assistance and use of facilities of the Laboratory for Artefact Studies, Faculty of Archaeology, Leiden University, the Netherlands. R.J. Looman (RMO) made the photographs of the barbed points. K. Lönnguist for assistance.

The collectors of the barbed points: Aad Berkhout, Rick van Bragt, Maarten Drummen, Gideon de Jong, Mirjam Kruizinga, Trudy Langeveld, Erwin van der Lee, Sibo van Maren, Johan Passchier, Maarten Schoemaker, Peter Soeters and Willy van Wingerden.

Contact: jan.dekker@york.ac.uk

Fig. 1. The analysed samples originate from a tight cluster of sites with no apparent tie between site and taxonomic identification.



Fig. 2 Schematic overview ZooMS workflow Amino acid string



### Conclusion

The presence of only Cervus elaphus/Alces alces and humans may be evidence for the preferential selection of particular species for barbed point production. Especially the use of human bone for projectile points is rare, but not unprecedented. A more widespread application of ZooMS will reveal whether this pattern is robust or the product of chance.



#### Results

Of the 10 barbed points nine provided results for ZooMS, isotopic analysis and <sup>14</sup>C dating (Fig. 1 & 3). Two species groups were identified using ZooMS: Cervus elaphus (red deer) /Alces alces (elk) and Homo sapiens (human) Fig. 4 shows the characteristic differences between the mass spectra of the two groups.

The  $\delta^{13}$ C and  $\delta^{15}$ N values of the two human points match with the values for Mesolithic human remains from the North Sea and the points identified as Cervus elaphus/Alces alces match with the range of variation of terrestrial herbivores.

All barbed points were dated to the Mesolithic with their dates ranging from 9505-7315 BP (uncal) or 11075-8021 BP (OxCal v4.3).



#### Fig. 3 $\delta^{13}$ C and $\delta^{15}$ N values of the barbed points

Fig. 4 Mass spectra of a red deer/elk (P06) and human (P29) barbed point

